

PCT


WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C07C 317/06, 321/06, 211/07, 49/84, C11D 1/72 // A61K 7/00, 9/107, B01F 17/00	A1	(11) International Publication Number: WO 98/37062 (43) International Publication Date: 27 August 1998 (27.08.98)
(21) International Application Number: PCT/US98/03103 (22) International Filing Date: 20 February 1998 (20.02.98) (30) Priority Data: 08/804,707 21 February 1997 (21.02.97) US (71) Applicant: RHONE-POULENC INC. [US/US]; CN-7500, Prospect Plains Road, Cranbury, NJ 08512-7500 (US). (72) Inventors: TRACY, David, James; 37-03 Ravens Crest Drive, Plainsboro, NJ 08536 (US). LI, Ruoxin; 1415 Ravens Crest Drive, Plainsboro, NJ 08536 (US). (74) Agents: SHEDDEN, John, A. et al.; Rhône-Poulenc Inc., Prospect Plains Road, Cranbury, NJ 08512-7500 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>

(54) Title: NOVEL NONYLPHENOL NONIONIC GEMINI SURFACTANTS**(57) Abstract**

Novel nonylphenol nonionic gemini surfactants are extremely effective emulsifiers for oil-in-water emulsions that provide improved detergency at even low concentration levels and are characterized by formula (I), wherein R and R₁ are both either H or a C₄-C₂ alkyl, aryl, or alkylaryl with the stipulation that they cannot both be H at the same time; and R₂ is independently -COR₃CO-, -CH₂N(R₄)CH₂-, CH₂N(R₅)R₆(R₅)NCH₂-, or R₇-D-R₇ wherein R₃ independently represents a C₁-C₆ alkyl, branched alkyl or aryl; R₄ represents a C₁ to C₄ alkyl, aryl, alkylaryl, hydroxy alkyl, haloalkyl or H, R₅ represents a C₁ to C₁₀ alkyl or aryl and R₆ independently represents a C₁ to C₁₀ alkyl, aryl, or alkylaryl, R₇ is a C₂ to C₄ alkyl and D independently represents -S-, -SO- and SO₂ with the stipulation that R₅ and R₆ can form a heterocyclic ring; EO represents ethylene oxy-radicals, x and y are whole numbers and x+y equals a number from 2 to 200 inclusive.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

NOVEL NONYLPHENOL NONIONIC GEMINI SURFACTANTS

This invention relates to improved surfactants useful as emulsifiers and in detergents and personal care products at very low concentrations. The surfactants also exhibit little to no toxicity which makes them useful in a wide
5 variety of applications including personal care, cosmetics, and pharmaceuticals.

Emulsification is an extremely important technology and it is a process which is used in detergency, emulsion polymerization, cosmetics, food, agrochemicals, paints, paper, transportation of crude oil, etc. Emulsifiers function as essential ingredients in personal care and household products; industrial and
10 institutional cleaners including hair shampoos, car washes, carpet shampoo, hand dishwashing liquids, latex foaming, oil recovery compositions; and the like.

In order to form a relatively stable emulsion, an emulsifier is required to adsorb at an oil-water interface to prevent emulsion droplet coalescence. The majority of emulsifiers are synthetic surfactants or natural products with
15 amphiphilic character. Presently, usage levels of surfactants for effective emulsification are usually above 0.1% active based on the total weight of the detergent solution which is used in the final use composition. For a given emulsifier system, it would be advantageous to use a lower amount of surfactant to reduce the cost and amount of surfactant waste discharged into the
20 environment; and to improve the performance of final products (e.g., the film forming and water resistance will be improved in latex paints and skin irritation will be reduced for cosmetic products).

While conventional surfactants generally have one hydrophilic group and one hydrophobic group, recently a class of compounds having at least two
25 hydrophobic groups and at least two hydrophilic groups have been introduced.

These have become known as "gemini surfactants" in the literature (Chemtech,
March 1993, pp 30 - 33), and J. American Chemical Soc., 115, 10083-10090,
(1993) and the references cited therein). Other gemini surfactant compounds,
that is, compounds having at least two hydrophilic groups and at least two
5 hydrophobic groups are also disclosed in literature but often are not referred to
expressly as gemini surfactants.

An intensive study of gemini surfactants as emulsifiers has been
made. From these studies, it has been found that gemini surfactants are
unexpectedly effective emulsifiers when used at very low concentrations in
10 comparison to conventional surfactants. This finding is reflected in superior
detergency at very low concentrations. Other performance properties related to
emulsification as mentioned above can be also improved using low
concentrations of gemini surfactants.

15

SUMMARY OF THE INVENTION

In accordance with the present invention, it has been discovered that
gemini surfactants of the type described and claimed herein can be extremely
effective emulsifiers for oil-in-water emulsions at lower concentrations than that
needed utilizing corresponding conventional surfactants. Furthermore, the
20 detergency is also improved over that provided by conventional surfactants at
these low concentration levels. These novel gemini surfactants are also able to
lower the monomer concentration of anionic surfactants when the two are mixed
in a blend. This results in less skin irritation and the other toxic side effects of
anionic surfactants.

25

3
DETAILED DESCRIPTION OF THE INVENTION

The novel surfactant compounds of the invention are based on certain gemini surfactants. As used herein, the term "gemini surfactants" is intended to mean surfactants having at least 2 hydrophobic groups and at least 2 hydrophilic groups per molecule.

A number of gemini surfactants are reported in the literature, see for example, Okahara et al., J. Japan Oil Chem. Soc. 746 (Yukagaku) (1989); Zhu et al., 67 JAOCS 7,459 (July 1990); Zhu et al., 68 JAOCS 7,539 (1991); Menger et al., J. Am. Chemical Soc. 113, 1451 (1991); Masuyama et al., 41 J. Japan Chem. Soc. 4,301 (1992); Zhu et al., 69 JAOCS 1,30 (Jan. 1992); Zhu et al., 69 JAOCS 7,626 July 1992); Menger et al., 115 J. Am. Chem. Soc. 2, 10083 (1993); Rosen, Chemtech 30 (March 1993); and Gao et al., 71 JAOCS 7,771 (July 1994), all of this literature incorporated herein by reference.

United States Patent No. 5,585,516 to Varadaraj et. al. discloses two tail- two head and two tail-one head surfactants including biphenolic hydrocarbon moieties. Also, gemini surfactants are briefly disclosed in U.S. Patent No. 2,374,354, to Kaplan; U.S. Nos. 2,524,218, and 2,530,147 to Bersworth (two hydrophobic tails and three hydrophilic heads); U.S. No. 3,244,724 to Guttman; and U.S. No. 5,160,450 to Okahara, et al., all of which are incorporated herein by reference.

The gemini surfactants of the present invention are nonylphenol, nonionic surfactants. The hydrophilic and hydrophobic groups of each surfactant moiety may be any of those known in conventional surfactants having one hydrophilic group and one hydrophobic group. For example, a typical nonionic gemini surfactant, e.g., a bis-polyoxyethylene alkyl ether, would contain two

4


polyoxyethylene alkyl ether moieties. However, non-ionic gemini surfactants prior to this are extremely rare.

Gemini surfactants are relatively quite new and few species have been reported or disclosed in the prior art. United States Patent No. 5,534,197 to Scheibel teaches a method for the preparation of a nonionic gemini surfactant wherein the hydrophilic head is a sugar or carbohydrate while the hydrophobic head is a long chain alkyl, the two joined by a short alkyl chain. U.S. Patent Nos. 3,888,797 and 3,855,156, both to Marumo, disclose a number of nonionic gemini surfactant species in which the hydrophobic portion is comprised of a long chain lower alkyl or alkene while the hydrophilic portion is comprised of an ethoxylate group. U.S. Patent No. 4,892,806 to Briggs and EP 0,688,781A1 to Adams also disclose sugar-based hydrophilic heads joined to the hydrophobic counterpart by a short chain carbon bridge.

Each moiety would contain a hydrophilic group, e.g., polyoxyethylene, and a hydrophobic group, e.g., an alkyl chain.

Nonionic gemini surfactants which are useful in the present invention include those of the formula:

I.



5


wherein both R and R₁ are the same and can independently be a C₄ to C₂₂ alkyl or hydrogen with the one caveat that both R and R₁ cannot be H; R₂ can independently be -COR₃CO-, -CH₂N(R₄)CH₂- and -CH₂N(R₅)R₆(R₅)NCH₂- or -
5 CH₂CH₂SO₂CH₂CH₂- or R₇-D-R₇ wherein R₃ independently represents a C₁ to C₆ alkyl, branched alkyl, or aryl, R₄ represents a C₁ to C₄ alkyl, aryl, alkylaryl, hydroxyalkyl, haloalkyl, or H; R₅ represents a C₁ to C₁₀ alkyl or aryl and R₆ represents C₁ to C₁₀ alkyl, aryl or alkylaryl, R₇ is a C₁ to C₄ alkyl, D is -S-, -SO-, -
10 SO₂- and R₅ and R₆ can independently form a heterocyclic ring, EO represents ethyleneoxy radicals and x and y are whole number integers with the proviso that x + y is a number from 2 to 200, inclusive.

With respect to the formula comprising the compounds of the present invention, the term "alkyl" includes substituted alkyl, especially the hydroxy-substituted derivatives thereof.

15 Preferably, the nonionic gemini surfactants of the present invention are selected from the following structures.

6


II.



wherein R, R₁, and R₃ have been hereinbefore defined and x+y is a whole number from 2-200 inclusive.

7


III.



wherein R and R₁ have been hereinbefore defined and x+y is a whole number from 2-200 inclusive.

5


IV.




wherein x+y = 20

8

V.



VI.




5

wherein x and y are whole numbers and $x+y = 2-200$ inclusive.

9
The compounds of the present invention can be made by the Lewis-Acid catalyzed addition of a dicarboxylic acid or acid chloride to an alkylphenol. For example, the addition of succinic acid chloride to nonylphenol may be represented by the following reaction.

5




Other diacids which would be employed comprise terephthalic acid, maleic acid, malonic acid, fumaric, succinic, glutaric, itaconic 2,2-dimethylglutaric, 2,4-dimethylglutaric, pimelic, cyclohexanedicarboxylic, and adipic acid.

10

Mannich reactions are those in which an active methylene compound is reacted with formaldehyde and ammonia or primary and secondary amines to yield β-amino-carbonyl compounds. Mannich reactions can be used to prepare amine-coupled products. An example would be the reaction of N, N'-dimethylethylene diamine and formaldehyde with nonylphenol.

10



Typical amines useful in the reaction include N, N'- dimethyl-1, 6-
 hexanediamine, methylamine, N,N'-dimethyl-1,3-propanediamine, ethylamine,
 propylamine, butylamine, piperazine, β-hydroxyethylamine, β-
 5 chloroethylamine, benzylamine, β-phenylethylamine and mixtures thereof.

This class of gemini surfactants as disclosed in the present invention can
 be used in providing improved emulsions which are operable at lower
 concentrations than the known surfactants of the prior art.

Typical useful phenolics are p-octylphenol, p-nonylphenol, p-
 10 dodecylphenol, p-(1,1,3,3-tetramethylbutyl)phenol, p-2,4-diamylphenol, 2,4-
 di(1,1,3,3-tetramethylbutyl) phenol, 2,4-ditertbutylphenol, 2-methyl-4-octylphenol,

2-methyl-4-nonylphenol, 2-t-butyl-4-nonylphenol, 2-acetyl-4-nonylphenol, 2-acetyl, 6-methyl-4-nonylphenol and mixtures thereof.

Where desired, the surfactants used in the present invention can be oxyalkylated by reacting the product with an alkylene oxide according to known methods, preferably in the presence of an alkaline catalyst.

For alkylation conditions and commonly used alkylating agents, see Amphoteric Surfactants Vol. 12, Ed. B. R. Bluestein and C. L. Hilton, Surfactant Science Series 1982, pg. 17 and references cited therein, the disclosures of which are incorporated herein by reference.

The surfactant compositions comprising the gemini nonionic compounds of the invention are extremely effective in aqueous solution at low concentrations as defined herein. The surfactants of the invention can be used in any amount needed for a particular application which can be easily determined by a skilled artisan without undue experimentation.

While the gemini surfactants of the invention can be used alone, it has been unexpectedly found that blends of the compounds of the invention with certain other conventional well known anionic, nonionic, cationic and amphoteric surfactants as well as polymers and enzymes provide synergistic effects in relation to critical micelle concentration (cmc) and surface tension reducing ability, and hence improved detergency.

Another important unexpected property of the Gemini Surfactants is their ability to significantly lower the ionic monomer concentration in mixtures of anionic/nonionic or anionic/nonionic/amphoteric or anionic/nonionic/cationic surfactant mixtures. It is well known that anionic monomers that are useful in personal care systems are responsible for higher toxicity and skin irritancy.

12

Further, anionic surfactant monomers are responsible for the deactivation of enzymes used in detergent systems and moreover, being anionic, they interact strongly with proteins and enzymes thereby deactivating them. Also anionic surfactants very often absorb onto polymers used in detergents, wastewater treatment systems and conditioning agents used in personal care applications and so on, thereby changing the charged nature of such polymers. This results in the complete loss or poor performance for the polymers in their application.

The use of gemini surfactants in very small amounts in the presence of an anionic surfactant such as lauryl ether sulfate (ESY) dramatically reduces the anionic monomers. With the addition of less than a 0.1 mole fraction of the gemini surfactant, the anionic concentration in the mixture is brought down by greater than an order of magnitude of ten. This dramatic reduction of the anionic monomer at these very low concentrations will result in the elimination of any detrimental effects discussed above and brought about as a result of the anionic monomers. Therefore, this ability to decrease the anionic (or cationic) monomers in solution should provide performance enhancement for the surfactant or the surfactant polymer/enzyme mixtures and the other performance additives very often found in detergents, and personal care formulations.


The following examples are provided to more fully disclose, describe and set forth the elements and embodiments of the surfactants of the present invention. They are for illustrative purposes only, and it is recognized that minor changes or alterations may be made to the reaction parameters, reactants and isolation procedures that are not disclosed herein. However, to the extent that any such changes do not materially alter the final product or result, such

13

alternative measures are deemed as falling within the spirit and scope of the invention as recited by the claims that follow.

A. Preparation of Octylphenol Gemini Structure IV

5 IV.



where $x + y = 20$.

Example 1 - Coupling


10 Aluminum chloride anhydrous, 30 g (0.2M), was added to 41.2 g (0.2 M) 4(1,1,3,3-tetramethylbutyl) phenol in 100 ml nitrobenzene at 5-10° C over a one hour period. A solution of succinyl chloride (15.6 g, 0.1 mole) in 50 ml nitrobenzene was added over a 30 minute period and the reaction was allowed to warm to room temperature and stirred overnight. The reaction yielded 22.0 g of
15 product.

14

Example 2 - Ethoxylate Preparation

Fifty (50) grams (0.1M) of the product of Example 1 was charged in an autoclave and to this was added 0.05 g potassium hydroxide flakes. The autoclave was heated to 120°C with a nitrogen sparge to remove water. The reactor was degassed by evacuating to 20" vacuum and releasing with nitrogen. After three cycles ethylene oxide was added (89 g., 2.0 M) at a temperature of 160°C and 60 psig. After addition was complete the reaction was cooled to 120°C, sparged with nitrogen and neutralized with acetic acid.


The residual product whose structure is set forth below as structure IV exhibited excellent surface activity as measured by surface tension measurement techniques.



$$x + y = 20$$

15

16

B. Preparation of Methylaminomethylenebis (4,6-Diamylphenol)**Intermediate A**

5

Example 3

2,4-diamylphenol (468 gm., 2.0M) was added to one mole of methylamine (40% in water) followed by two moles of formaldehyde (as formalin). The temperature was kept at 25° to 35°C during the addition. The reaction was stirred overnight. The organic and aqueous phases were separated and the organic phase yielded the bisphenolic compound (Intermediate A) whose composition appeared as a yellow oil.

10

Example 4 - Preparation of the Ethoxylate of Intermediate A.

To an autoclave was charged one mole of the intermediate bisphenolic A together with 5 g of potassium hydroxide flakes. The autoclave was heated to 140°C and purged with nitrogen to remove water. The reactor was heated to 160°C and ethylene oxide added to a pressure of 60 psig until 20 moles had reacted. The reactor was cooled to 120°C, the product neutralized with acetic acid and discharged. The compounds' NMR analysis spectrum agreed with the predicted structure. The product foamed when added to water and exhibited


15

20

superior surface activity when measured by standard surface tension measurement techniques.

Example 5 - Preparation of Sulfone

5




4(1,1,3,3-tetramethylbutyl)phenol (87 g, 0.42 mol) and vinyl sulfone (10 g, 0.086 M) were dissolved in 100 ml dry xylene at room temperature. The reaction solution was refluxed for 8 hours in the presence of trace pure acetic acid as catalyst. The reaction was then stopped by stripping out all the solvent and excess starting material under vacuum. The final product was collected after cooling. The reaction yielded about 40 g of product.

15 **Example 6 - Ethoxylation of Example 5**

To an autoclave was charged 53 g (0.1 mole) of the product of Example 5 and 0.1 g of potassium hydroxide flakes were added. The autoclave was heated to 120°C with a nitrogen sparge to remove water. After degassing, ethylene oxide (3.0 mole, 132 g) was added at 150°C to 160°C and 40 psig. After uptake


was complete, the reaction was cooled to 110°C and neutralized with glacial acetic acid. The residue exhibited surface activity as evidenced by surface tension reduction and foaming characteristics.



18

What We Claim Is:

- 1) A surfactant composition comprising one or more of the compounds comprised of the formula:




5

- wherein R and R₁ are both either H or a C₄-C₂₂ alkyl, aryl, or alkylaryl with the stipulation that they cannot both be H at the same time; and R₂ is independently -COR₃CO-, -CH₂N(R₄)CH₂-, CH₂N(R₅)R₆(R₅)NCH₂-, or R₇-D-R₇ wherein R₃ independently represents a C₁-C₆ alkyl, branched alkyl or aryl; R₄ represents a C₁ to C₄ alkyl, aryl, alkylaryl, hydroxy alkyl, haloalkyl or H, R₅ represents a C₁ to C₁₀ alkyl or aryl and R₆ independently represents a C₁ to C₁₀ alkyl, aryl, or alkylaryl, R₇ is a C₂ to C₄ alkyl and D independently represents -S-, -SO- and -SO₂- with the stipulation that R₅ and R₆ can form a heterocyclic ring; EO represents ethylene oxy-radicals, x and y are whole numbers and x+y equals a number from 2 to 200 inclusive.

19


- 2) The composition of claim 1 further characterized by the structure:



wherein R and R_1 have been hereinbefore defined and R_3 independently
5 represents a C_1 - C_6 alkyl, branched alkyl, or aryl and x and y are whole numbers
and $x+y$ equals a number from 2-200 inclusive.


20

- 3) The composition of claim 2 further characterized by the structure.



- 5 where R and R₁ have been hereinbefore defined and x and y are whole numbers and x+y equals a number from 2-200 inclusive.


- 21
4) The composition of claim 3 further characterized by the structure:



wherein $x+y$ are whole numbers

and $x+y = 20$

- 5) The composition of claim 1 further characterized by the structure:




wherein R and R₁ have been previously

defined and $x+y$ equals a whole number from 2-200 inclusive.

22

- 6) The composition of claim 5 further characterized by the structure:




and $x+y$ equals a number of from

5 2-200 inclusive.


23

- 7) The composition of claim 1 further characterized by the structure.



- 5 wherein R and R_1 have been herein before defined, x and y are whole numbers and $x+y$ equals a number from 2-200 inclusive.


8) The composition of claim 7 further characterized by the structure:



wherein x and y are whole numbers

and $x+y$ equals a number from 2-200 inclusive.

- 9) The composition of claim 1 ²⁵ further characterized by the structure:



wherein x and y are whole numbers and

$x+y =$ a number from 2-200 inclusive

5

- 10) A surfactant blend comprising the compound of claim 1 and one or more additional compounds selected from the group consisting of anionic surfactants, nonionic surfactants, cationic surfactants, amphoteric surfactants, and mixtures thereof.

10

- 11) A surfactant blend comprising the compound of claim 2 and one or more additional compounds selected from the group consisting of anionic surfactants,

26

nonionic surfactants, cationic surfactants, amphoteric surfactants, and mixtures thereof.

12) A surfactant blend comprising the compound of claim 3 and one or more
5 additional compounds selected from the group consisting of anionic surfactants,
nonionic surfactants, cationic surfactants, amphoteric surfactants, and mixtures
thereof.

13) A surfactant blend comprising the compound of claim 5 and one or more
10 additional compounds selected from the group consisting of anionic surfactants,
nonionic surfactants, cationic surfactants, amphoteric surfactants, and mixtures
thereof.

14) A surfactant blend comprising the compound of claim 7 and one or more
15 additional compounds selected from the group consisting of anionic surfactants,
nonionic surfactants, cationic surfactants, amphoteric surfactants, and mixtures
thereof.

15) A surfactant blend comprising the compound of claim 9 and one or more
20 additional compounds selected from the group consisting of anionic surfactants,
nonionic surfactants, cationic surfactants, amphoteric surfactants, and mixtures
thereof.

- 27
- 16) The surfactant composition of claim 1 useful as an emulsifier in detergents, paints, cosmetics, foods, shampoos, oil recovery compositions, dishwashing liquids, and latex foaming compositions.
- 5 17) The surfactant composition of claim 3 useful as an emulsifier in detergents, paints, cosmetics, foods, shampoos, oil recovery compositions, dishwashing liquids, and latex foaming compositions.
- 10 18) The surfactant composition of claim 5 useful as an emulsifier in detergents, paints, cosmetics, foods, shampoos, oil recovery compositions, dishwashing liquids, and latex foaming compositions.
- 15 19) The surfactant composition of claim 7 useful as an emulsifier in detergents, paints, cosmetics, foods, shampoos, oil recovery compositions, dishwashing liquids, and latex foaming compositions.
- 20) The surfactant composition of claim 9 useful as an emulsifier in detergents, paints, cosmetics, foods, shampoos, oil recovery compositions, dishwashing liquids, and latex foaming compositions.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 98/03103

A. CLASSIFICATION OF SUBJECT MATTER

IPC6: C07C 317/06, C07C 321/06, C07C 211/07, C07C 49/84, C11D 1/72 // A61K 7/00,
A61K 9/107, B01F 17/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: C11D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WPI, CA, CLAIMS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	US 5710121 A (DAVID JAMES TRACY ET AL), 20 January 1998 (20.01.98), Corresponding WO 97/23449; see the whole document --	1-4, 10-12, 16-17
X	US 5493050 A (RAMESH VARADARAJ ET AL), 20 February 1996 (20.02.96) --	1
X	US 5585516 A (RAMESH VARADARAJ ET AL), 17 December 1996 (17.12.96) --	1
X	GB 2150569 A (SANDOZ LTD), 3 July 1985 (03.07.85) --	1

☒ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

- * Special categories of cited documents:
- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search

30 June 1998

Date of mailing of the international search report

10 AUG 1998

Name and mailing address of the ISA:



European Patent Office, P.B. 5818 Patentlaan 2
NL-2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

DAGMAR JÄRVMAN

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 98/03103

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 4592809 A (JAAN J. FONG ET AL), 3 June 1986 (03.06.86) --	1
A	US 3919429 A (MAX GROSSMANN ET AL), 11 November 1975 (11.11.75) -- -----	1

INTERNATIONAL SEARCH REPORT
Information on patent family members

09/06/98

International application No.

PCT/US 98/03103

Patent document cited in search report			Publication date	Patent family member(s)		Publication date	
US	5710121	A	20/01/98	AU	1069597 A	17/07/97	
				WO	9723449 A	03/07/97	

US	5493050	A	20/02/96	NONE			

US	5585516	A	17/12/96	CA	2170937 A	11/09/96	
				EP	0732323 A	18/09/96	

GB	2150569	A	03/07/85	CH	666480 A,B	29/07/88	
				DE	3442888 A,C	13/06/85	
				FR	2555986 A,B	07/06/85	
				HK	191 A	11/01/91	
				JP	1776985 C	28/07/93	
				JP	4061865 B	02/10/92	
				JP	60132947 A	16/07/85	
				US	5470355 A	28/11/95	

US	4592809	A	03/06/86	AU	5297886 A	12/02/87	
				CA	1268184 A	24/04/90	
				DE	3611627 A	19/02/87	
				GB	2178747 A,B	18/02/87	
				JP	62033147 A	13/02/87	

US	3919429	A	11/11/75	AU	5112873 A	18/07/74	
				BE	794214 A	18/07/73	
				DD	104020 A	20/02/74	
				DE	2300860 A,B,C	23/08/73	
				FR	2168474 A,B	31/08/73	
				GB	1388251 A	26/03/75	
				JP	48079894 A	26/10/73	
				NL	7300488 A	20/07/73	
				ZA	7300297 A	31/10/73	

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☒ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.